
2012 KENTUCKY ASTHMA PROGRAM FACT SHEET: PURCHASE AREA DEVELOPMENT DISTRICT


- The Purchase District of Kentucky is comprised of 8 counties: Ballard, Carlisle, Hickman, Fulton, McCracken, Graves, Marshall, and Calloway.
- The 2011 Purchase District population was roughly 197,000. About 42,300 persons were younger than 18 (children), and about 154,800 were 18 or older (adults).



PREVALENCE

In 2010, an estimated 6,500 children (15.8%) in the Purchase District currently had asthma, compared to 10.7% for Kentucky and 8.4% for the United States. In 2011, an estimated 12,100 adults (7.8%) in this district currently had asthma, compared to 10.4% for Kentucky and 9.1% for the United States.







In 2010, child current asthma prevalence rate was higher in boys as compared to Kentucky. In 2011, adult current asthma prevalence rate was lower in both males and females in this district compared to Kentucky. (NOTE: BRFSS 2010 data cannot be compared to BRFSS 2011 data because of change in methodology)

HOSPITALIZATIONS

In 2011, the age-adjusted hospitalization rate for children was 6.9/10,000 persons compared to the Kentucky rate of 4.9/10,000 persons; while the age-adjusted hospitalization rate for adults was 8.2/10,000 persons compared to the Kentucky rate of 9.7/10,000 persons.





In 2011, child asthma hospitalizations were higher in Purchase District compared to the state, while adult asthma hospitalizations in this district were lower compared to the state. Between 2001 and 2011, this district averaged 11.7 asthma hospitalizations/10,000 persons/year, while Kentucky averaged 8.3 asthma hospitalizations/10,000 persons/year over the same 11 year period.²

EMERGENCY DEPARTMENT VISITS

In 2011, the asthma Emergency Department (ED) visits for children were 1.7/1,000 persons, compared to the state rate of 1.9/1,000 persons³ and for adults were 2.1/1,000 persons compared to the state rate of 2.5/1,000 persons.³

MORTALITY

Between 2001 and 2010, the average age-adjusted asthma deaths were 0.91/100,000 persons/year in this district compared to 1.10/100,000 persons/year for Kentucky. In 2010, crude asthma death rates for Kentucky were 1.01/100,000 persons⁴ compared to the US crude death rate of 1.1/100,000 persons.⁵

NOTES:

- 1. Behavioral Risk Factor Surveillance System (BRFSS).
- 3. OHP: Preliminary data 2011.

- 2. Kentucky. Office of Health Policy (OHP), 2001-2011.
- 4. Office of Vital Statistics, Kentucky, 2001-2010.

5. CDC Wonder data: http://wonder.cdc.gov/ucd-icd10.html This fact sheet was supported by the Cooperative Agreement Number 5U59EH000508-04 from the Centers for Disease Control and Prevention (CDC). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the CDC.

KENTUCKY RESPIRATORY DISEASES ASTHMA PROGRAM

Jan Hatfield, Program Manager; Sue Thomas-Cox, PI; Deepa Valvi, Asthma Epidemiologist 275 E. Main St. * Mail Stop HS2WE * Frankfort, KY 40621 * (502) 564-7996, ext. 4439

Website: www.chfs.ky.gov/Asthma